




Ohio State Fact Sheet

Leading Causes of death in Ohio in 2013*



Ohio has the 15th highest death rate from cardiovascular disease in the country.**

- Heart disease is the No. 1 killer in Ohio*
- Stroke is the No. 4 killer in Ohio*
- 26,878 people in Ohio died of heart disease in 2013*
- 5,690 people in Ohio died of stroke in 2013*

Heart Disease and Stroke Risk Factors in Ohio

	Ohio	US
Adults who are current smokers	23.4%	19%
Adults who participate in 150+ min of aerobic physical activity per week	49.5%	50.5%
Adults who are overweight or obese+	65.1%	64.3%
Adults who have been told that they have had a heart attack	5.3%	4.4%
Adults who have been told that they have had a stroke	3.7%	2.8%
Adults who have been told that they have angina or coronary heart disease	4.7%	4.1%
Population of adults (18-64) who have some kind of health care coverage	83.3%	80%
High school Students who are obese++	13%	13.7%

* List includes Puerto Rico and D.C. Based on total number of deaths in 2013. Centers for Disease Control and Prevention. WISQARS Leading Cause of Death Reports, 2013.
 ^ Also known as COPD (Chronic Obstructive Pulmonary Disorder); COPD and allied conditions (including asthma); the term in the ICD/10 is "chronic lower respiratory diseases."
 ** Based on 2010 age-adjusted death rates. American Heart Association. Heart Disease and Stroke Statistics: 2015 Update. A Report from the American Heart Association. Circulation, Accessed February 1, 2015
 ***Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System Survey, 2013.
 + Overweight is defined as having a body mass index (BMI) of 25.0-29.9. Obese is defined as having a body mass index of 30.0 or more.
 ++ Students who were ≥ 95th percentile for body mass index, by age and sex. Centers for Disease Control and Prevention, Youth Risk Behavior Surveillance System, 2013.